

not just a  
'cry for  
help'



prevention of young suicide

This booklet is for you  
if you know someone  
who has made a  
suicide attempt.



It gives some guidance on  
how you can help him or her  
through this crisis.

Knowing that someone we  
care about is feeling suicidal  
can be emotionally and physically  
draining. It is important to look  
after your own health and to  
make time to get support and  
advice for yourself too.

# You are not alone

Attempted suicide is far more common than most of us realise. In England and Wales, thousands of people are admitted to accident and emergency departments each year because they have tried to kill themselves.

Although most people who attempt suicide survive, a few remain at high risk of taking their own life for quite some time afterwards. This means that any suicide attempt, however minor it seems to be, should be taken seriously.

## Why did it happen?

### Thinking about suicide

Suicidal thoughts can coincide with times of change – whether these changes are ‘good’ (like getting married or starting a new job), or ‘bad’ (such as someone dying or the end of a relationship). Quite often two or three different things will have happened, each of them causing considerable stress.

For some people there may be no apparent reason for these thoughts.

The circumstances leading up to a suicide attempt are different for everybody, and the reasons for it happening may never be fully explained or understood.

Often those who think about suicide are depressed. Usually very sensitive people, they may have been in despair, feeling hopeless about the future and unable to think straight. Everyday worries have become totally out of proportion, and black thoughts may have left no room for anyone or anything else.

## Acting on the thought

Turning thoughts about suicide into action is sometimes done on impulse, perhaps following an event that is seen as 'the last straw'. This is especially true of young people.

Others, however, may have been making plans about suicide for some time.

Using alcohol and/or other drugs can make things worse. They take away the inhibitions which would otherwise stop someone from attempting to take their own life.

## It's hard to talk ...

... about fears and feelings – even to those we know love and care about us. This can prevent other people from recognising the distress and being able to help in a crisis.

Words are often totally inadequate to convey the amount of pain a person may be suffering.

It is easy to understand that someone is hurting if they have been badly injured or are physically ill. Emotional pain cannot be seen, but it can be just as unbearable.

Sometimes attempting suicide may be the only way to show other people how bad things are.

When your back is up against the wall, suicide can seem to be the only way out.



# What can be done to help?

## ...stay ALERT

**Ask** them how they were feeling before it happened and how they are feeling now.

Talking about suicide does not make it more likely to happen. Try to be patient if they are angry or refuse to talk. It may be that writing things down is an easier way for them to communicate with you.

**Listen** – this is the most important thing you can do. Treat them with respect, and try not to be judgmental or critical.

**Empathise** by showing that you really are trying to understand things from their point of view.

Words don't always matter. The touch of a hand or a hug can go a long way to show that you care.

**Reassure** them that desperate feelings are very common and can be overcome.

Things can and do change, help can be found and there is hope for the future. People DO get better!

**Try** to give practical support.

Help them to cope with any extra pressures. It may not be possible to deal with all the things that are troubling them, but between you agree on what you will do if a suicidal crisis happens again.

# ...don't

**Put** them down or do things that might make them feel worse. A suicide attempt suggests that self-esteem is already very low.

**Abandon** or reject them in any way. Your help, support and attention are vital if they are to begin to feel that life is worth living again. Don't relax your attentions just because they seem to be better. It doesn't mean that life is back to normal for them yet. They may be at risk for quite a while.

**Nag** – although it may be well meant. Nobody wants to be pestered all the time. Don't intrude – try to balance being watchful with a respect for privacy.

**Ignore** what has happened.

**Criticise** their actions – however you may be feeling about their suicide attempt, try to remember the pain and turmoil that they were, and may still be, going through. Don't take their behaviour personally – it was not necessarily directed at you.

# Help from services

There are many ways in which health workers, social services staff and others can help someone who has attempted suicide. The staff in the hospital accident and emergency department may make a referral, or the family doctor can be asked to.

## Treating depression:

Suicidal feelings may be linked to being depressed. Depression is a serious illness but can be successfully treated by anti-depressant medication and/or 'talking and listening treatments'.

## Extra help in a crisis

If you feel that the situation is getting worse rather than better, and you are worried about another suicide attempt, trust your instinct and share your concerns straight away.

Contact the family doctor, or any professional who has been involved already.

These people cannot discuss confidential details with you, but they will want to listen to what you are saying and **they can follow it up.**

...or for further help  
phone the PAPHYRUS helpline  
**HOPELineUK**  
(see overleaf)...



Call **HOPELineUK**

0800 068 41 41

**HOPELineUK** is a national confidential telephone service providing support, practical advice and information to anyone concerned that a young person they know may be at risk of suicide.

**HOPELineUK** is staffed by trained professionals who can discuss your needs in coping with someone who is feeling suicidal.

Our advisors have access to a comprehensive database from which they can give you contact details of further sources of help.

The helpline is open on weekdays from 10am – 5pm, and 7pm – 10pm and on weekends from 2pm – 5pm.

Calls are free from BT landlines. If you call during busy times or when the helpline is closed you can leave your contact details and an advisor will call you back.



Lodge House, Thompson Park, Burnley BB11 2RU.

**Tel:** 01282 432555

**e-mail:** [admin@papyrus-uk.org](mailto:admin@papyrus-uk.org)

**[www.papyrus-uk.org](http://www.papyrus-uk.org)**

This resource is funded by the  
**Department for Children, Schools and Families.**

© PAPYRUS 2008

Company Limited by Guarantee Number 3555482  
Registered office as above. Registered Charity Number 1070896

---

This booklet is intended for guidance only and is not a substitute for professional advice, no responsibility for loss occasioned as a result of any person acting or refraining from acting because of what is written in the booklet can be accepted by the publisher or authors or PAPYRUS's Trustees.